

その他出版物（英文）

- 1) Inui S, Okada N, Yoshikawa K, Harino S: Von Recklinghausen's disease in monozygotic twins. Proceedings: The 8th Japan-Korea Joint Meeting of Dermatology, 589-591,1993.
- 2) Inui S, Fukuzato Y, Nakajima T, Yoshikawa K, Itami S: Identification of androgen-inducible TGF- β 1 derived from dermal papilla cells as a key mediator in androgenetic alopecia. J Invest Derm Symp Proc 8: 69-71, 2003.
- 3) Itami S, Inui S: Role of androgen in mesenchymal epithelial interactions in human hair follicle. J Investig Dermatol Symp Proc 10: 209-211, 2005.
- 4) Kitagawa T, Matsuda K, Inui S, Takenaka H, Katoh N, Itami S, Kishimoto S, Kawata M: Keratinocyte growth inhibition through the modification of Wnt signaling by androgen in balding dermal papilla cells. Mol Endocrinol 23: 279, 2009. (Published as Translational Highlights from J Clin Endocrinol Metab) (4.746)
- 5) Imagawa K, Inui S: A visit to the house of Dr. Shoji Okuda. Hair Transplant Forum International 19: 200-201, 2009.
- 6) Inui S: *In vitro* and *in vivo* evidence of pathogenic roles of Hic-5/ARA55 in keloids through Smad pathway and profibrotic transcription. J Wound Tech, in press.
- 7) Inui S, Shono F, Nakajima T, Hosokawa K, Itami S: Identification and characterization of cartilage oligomeric matrix protein (COMP) as a novel pathogenic factor in keloids. J Wound Tech, in press.